

Expression analysis of heat-shock protein gene *Hsp845* in the Antarctic psychrotrophic bacterium *Psychrobacter* sp. G under temperature and salinity stress

CHE Shuai^{1,2}, ZHANG Liang^{1,2}, ZHANG Zheng^{1,2} & LIN Xuezheng^{1,2*}

¹ The First Institute of Oceanography, SOA, Qingdao 266061, China;

² Key Lab of Marine Bioactive Substances, SOA, Qingdao 266061, China

Received 4 January 2016; accepted 15 March 2016

Abstract Heat shock proteins (Hsps), produced by organisms under high temperature stimulation, play important roles in protein folding, translocation, and refolding/degradation. In this study, we investigated the expression level of the GrpE Hsp gene *Hsp845* of *Psychrobacter* sp. G under different temperature and salinity stresses by quantitative real-time PCR and western blotting, respectively. At both transcriptional and translational levels, *Hsp845* gene expression was induced by high temperature (30°C) and inhibited by low temperatures (0°C and 10°C). *Hsp845* expression was also induced both by the absence of salt (0‰) and high salinity (90‰ and 120‰) at the transcriptional level, but was only induced by high salinity (90‰ and 120‰) at the translational level. In a combined stress treatment, *Hsp845* was more sensitive to high temperature than to salinity at both transcriptional and translational levels. The increase in the translational-level expression of *Hsp845* lagged behind that at the transcriptional level, and *Hsp845* maximum expression was also higher at the transcriptional than at the translational level. In the absence of salt, transcriptional- and translational-level expressions exhibited opposite patterns, suggesting that the underlying mechanism requires further study.

Keywords *Psychrobacter*, *Hsp845*, GrpE, stress response, qRT-PCR, western blotting

Citation: Che S, Zhang L, Zhang Z, et al. Expression analysis of heat-shock protein gene *Hsp845* in the Antarctic psychrotrophic bacterium *Psychrobacter* sp. G under temperature and salinity stress. *Adv Polar Sci*, 27: 14-20, doi: 10.13679/j.advps.2016.1.00014

1 Introduction

Heat shock response, which involves the synthesis of heat shock proteins (Hsps), plays a vital role in the defense of bacterial cells against temperature-induced damage. Hsps are highly conserved and exist in almost all species^[1]. GrpE, a member of the Hsp family, is transiently induced under various stress conditions^[2]. GrpE is a known nucleotide-exchange factor for the molecular chaperone DnaK, which functions co-translationally and post-translationally to promote protein folding and disaggregation in cells^[3-6]. In

this molecular chaperone system, DnaK (an *Hsp70* homolog) acts in concert with GrpE and its co-chaperone DnaJ, an *Hsp40* homolog^[7]. DnaJ stimulates the hydrolysis of DnaK-bound ATP and converts DnaK from a low-affinity state to a high-affinity state for substrates, whereas GrpE facilitates ADP/ATP exchange and restores DnaK to the low-affinity state^[8-10]. GrpE gene expression is regulated by $\sigma 70$ and $\sigma 32$ promoters^[11], thereby allowing rapid and transient induction in response to elevated temperatures^[12-13]. Previous studies have also shown that this molecular chaperone system is essential to normal cell growth^[14-16], cell division^[16], membrane translocation^[17], chromosome segregation, and maintenance of low-copy-number plasmids^[3]. As Hsps, the

* Corresponding author, E-mail: linxz@fio.org.cn

expressions of DnaK, DnaJ, and GrpE are transiently induced under different stress conditions, such as heat^[18], salinity^[19], ethanol treatment, viral infection, and exposure to heavy metals^[14,20].

Sea ice in the Southern Ocean is dominated by strong gradients of salinity and drastic fluctuations of temperature, space, and light^[21]. Temperature and salinity are the two most important factors affecting the survival of microorganisms living in sea ice. These organisms have accordingly evolved adaptive mechanisms ranging from molecular to whole cell levels. The freezing and thawing of ice during different seasons in Antarctic seawater inhabited by *Psychrobacter* sp. affects the salinity of offshore seawater^[22].

Although the heat-shock response of GrpE is well documented at the molecular level in mesophilic microorganisms such as *Escherichia coli*^[12,14], *Synechocystis* sp.^[15], and *Bacillus subtilis*^[23] and in thermophilic microorganisms such as *Thermus thermophilus*^[24] and *Thermosynechococcus elongates*^[15], limited studies have been performed in cold-adapted microorganisms. Moreover, most of the earlier studies focused on response to only one stress factor, such as temperature. In fact, organisms in nature seldom undergo only one stress at a time^[1]. To our knowledge, researchers have rarely investigated the mechanism of GrpE in response to salinity and to combinations of multiple stresses at both transcriptional and translational levels simultaneously. According to our previous study^[2], the *Hsp845* gene is 615 bp long and encodes a protein consisting of 204 amino acid residues with a molecular mass of 22.7 kDa and a calculated isoelectric point of 4.54. A BLAST analysis revealed that the deduced amino acid sequence of *Hsp845* was a close homolog to amino acid sequences of other known cytosolic GrpEs. Among the four Hsp genes we investigated, *Hsp845* was found to be strongly sensitive at the transcriptional level to temperature and salinity stress^[2]. In the present study, we further investigated the expression of the *Hsp845* gene under different conditions (temperature, salinity, and combinations of temperature and salinity) at both transcriptional and translational levels by quantitative real-time PCR (qRT-PCR) and western blotting, respectively.

2 Materials and methods

2.1 Bacteria, medium, and growth conditions

The Antarctic psychrotrophic bacterium *Psychrobacter* sp. G was isolated from Antarctic seawater and maintained in our laboratory^[25]. *Psychrobacter* sp. G was cultured at 20°C with rotation of 150 r·min⁻¹ in medium consisting of 5 g tryptone, 1 g yeast extract and 14 g NaCl in 1 L seawater. The composition of the medium used for salinity stress treatments, which has been described previously, is summarized in Table 1^[26]. Seawater (with an approximate salinity of 31‰) was taken from the coast at Qingdao (Shandong, China).

Table 1 Components of culture medium having different salinities

Salinity	Tryptone /g	Yeast extract/g	NaCl /g	Seawater /mL	Deionized water/mL
0	5	1	0	0	1000
15	5	1	0	480	520
45	5	1	14	1000	0
90	5	1	59	1000	0
120	5	1	89	1000	0

2.2 Recombinant expression and purification of *Hsp845* protein

Primers *Hsp845F* (5'-GGAATTCATATGAGCGAGCAAAATAACCAC-3') and *Hsp845R* (5'-CCGCTCGAGCTGACCGACACGTACCATCG-3'), respectively containing NdeI and XhoI restriction enzyme sites, were designed to amplify the open reading frame of the *Hsp845* gene. After digestion with restriction enzymes NdeI and XhoI, the target PCR product was subcloned into plasmid pET22b digested with the same restriction enzymes. The recombinant plasmid was then transformed into *E. coli* BL21 (DE3) competent cells for recombinant protein expression. Positive clones were screened by PCR with primers T7F and T7R and then confirmed by nucleotide sequencing. Each positive clone was incubated in 250 mL Luria-Bertani medium (containing 100 µg·mL⁻¹ ampicillin) at 37°C with shaking at 150 r·min⁻¹. When the absorbance of the culture at 600 nm (OD₆₀₀) reached 0.6–0.8, the cells were incubated for an additional 4 h in the presence of the inducer isopropyl-β-D-thiogalactopyranoside (IPTG) added to a final concentration of 1 mM. The cells were then collected by centrifugation at 10000×g for 5 min and resuspended in 5 mL of 50 mM Tris-HCl buffer (pH 8.0). After lysis of the cells with a one-shot cell disrupter (UL61010A-1, Constant Systems Ltd., Daventry, UK), the lysates were centrifuged at 10000×g for 20 min at 4°C. The recombinant protein was purified on a Ni Sepharose 6 Fast Flow column (11-0008-86 AE, GE Healthcare, Uppsala, Sweden) according to the manufacturer's instructions. The antiserum was prepared and validated by GenScript (Nanjing, China; order ID: 168267-1).

2.3 Stress treatments

Psychrobacter sp. G was initially cultured at its optimal growth temperature (20°C) and salinity (45‰)^[26]. When the OD₆₀₀ of the culture reached approximately 0.5, temperature, salinity, and combined stress treatments were carried out as follows. For temperature stress treatments, cultures were kept at different temperatures (0°C, 10°C, or 30°C) for different durations (2 h, 6 h, or 12 h). For salinity stress treatments, the cultures were centrifuged at 8000 ×g for 5 min at 20°C. After thoroughly suspending the pellets with an equal volume of

medium to a final salinity of 0‰, 15‰, 90‰, or 120‰, the suspensions were maintained at 20°C for 2 h, 6 h, or 12 h. In the combined stress treatments, cells were cultured under the following conditions: a final salinity of 15‰ at 0°C or 30°C or a final salinity of 90‰ at 0°C or 30°C. A strain cultured continuously under optimal conditions was used as the control for qRT-PCR and western blot analysis. Three biological replicates of each control and stress treatment were carried out.

2.4 Total RNA isolation and cDNA synthesis

Total RNA was extracted using an RNAprep Pure Cell/Bacteria kit (DP430, Tiangen Biotech, Beijing, China) according to the manufacturer's protocol. The purity of prepared RNA was judged by the A_{260}/A_{280} ratio. A PrimeScript RT Reagent kit (DRR037A, Takara, Dalian, China) was used for the reverse transcription of RNA into cDNA. The cDNA for qRT-PCR was prepared using 500 ng of total RNA. Primers Q-F (5'-TGGTGGATAACTTGGA-3') and Q-R (5'-ACCGACCGCTTCAT-3') for qRT-PCR were designed with the software program Primer 5.0. The GAPDH gene encoding glyceraldehyde-3-phosphate dehydrogenase was used as an internal reference^[2,26].

2.5 Expression analysis of the *Hsp845* gene by qRT-PCR

SYBR *Premix Ex Taq* II (DRR820A, Takara) was used for the qRT-PCR analysis. qRT-PCR amplification was carried out on a Stratagene Mx3005P qPCR System for 40 cycles (95°C for 1 min, 45°C for 30 s, and 72°C for 20 s). All reactions were performed in triplicate. Quantification of mRNA was based on threshold cycle (C_t) values. After normalizing the C_t value of *Hsp845* using the C_t value corresponding to the selected reference gene, the efficiency of each qRT-PCR amplification was calculated. Data analysis was based on the comparative C_t ($2^{-\Delta\Delta C_t}$) method^[27]. Data obtained from qRT-PCR analysis were subjected to analysis of variance to identify significant differences between mean values among treatments. Differences were considered to be statistically significant at $P < 0.05$ and highly significant at $P < 0.01$.

2.6 Western blot analysis of *Hsp845* protein expression in *Psychrobacter* sp. G

Western blot analysis was carried out to investigate the expression of *Hsp845* at the translational level. Total protein of *Psychrobacter* sp. G was dissolved in 50 mM Tris-HCl buffer (pH 8.0) and then centrifuged at 10000×g for 10 min at 4°C to collect the supernatant. Protein concentration was then assayed. Each sample (approximately 60 mg protein) was analyzed by 15% sodium dodecyl sulfate-polyacrylamide gel electrophoresis. The proteins were then electroblotted onto Immobilon-P transfer membranes (IPVH00011, SolarBio,

Beijing, China). To prevent nonspecific antibody binding, membranes were first blocked with 5% non-fat milk in TBST (10 mM Tris-HCl [pH 7.5], 150 mM NaCl, and 0.1% Tween-20) accompanied by shaking for 3 h at room temperature and then incubated with the 1:1000 diluted antiserum to *Hsp845* in TBST overnight. After three 10-min washes with TBST and one 10-min wash with TBS (10 mM Tris-HCl [pH 7.5], 150 mM NaCl), each membrane was incubated in peroxidase-conjugated goat anti-rabbit IgG (1:2000 in block reagents as indicated above) for 2 h and then washed three times with TBST (10 min per wash) followed by a 10-min wash in TBS. Finally, the membrane was soaked in a reaction mixture (1 mL of 6 mg·mL⁻¹ 4-chloro-1-naphthol in methanol, 10 mL TBS, and 6 μL H₂O₂) in darkness for 10 min and the reaction was blocked in distilled water. Antibody binding was visualized through a peroxidase-catalyzed stain reaction^[28].

3 Results

3.1 Expression and purification of recombinant *Hsp845* protein

Recombinant *Hsp845* protein was expressed in *E. coli* BL21 (DE3) induced by IPTG (Figure 1, lanes 2 and 3). Expressed recombinant *Hsp845* protein carrying a 1.08-kDa His-tag was detected in the lysate supernatant by Coomassie Bright Blue staining and found to have a molecular mass of approximately 24 kDa, consistent with the expected value. The recombinant *Hsp845* protein was purified by Ni Sepharose 6 Fast Flow chromatography (Figure 1, lane 4), which revealed that the purity of the obtained protein was higher than the threshold criterion of the antibody test (85%). The recombinant protein was sent to GenScript to revalidate its purity and obtain effective antigen protein.




Figure 1 Sodium dodecyl sulfate-polyacrylamide gel electrophoretic analysis of recombinant expression of *Hsp845* protein. Lane 1, protein marker; lane 2, lysate of *Escherichia coli* with pET22b-*Hsp845* without induction; lane 3, lysate of *E. coli* with pET22b-*Hsp845* induced by isopropyl-β-D-thiogalactopyranoside; lane 4, recombinant *Hsp845* protein purified by Ni Sepharose 6 Fast Flow chromatography.

3.2 Expression characteristics of *Hsp845* in response to temperature stress

qRT-PCR analysis revealed that *Hsp845* gene expression was significantly inhibited by low temperatures (0°C and 10°C) and enhanced immediately after exposure to high temperature (30°C). Under 30°C conditions, *Hsp845* gene expression reached a maximum after 2 h of treatment, with a 5.3-fold increase compared with the control (20°C), and then decreased during the next 10 h (Figure 2). Western blot analysis indicated that the expression of *Hsp845* protein was also enhanced by high temperature (30°C) and inhibited by low temperatures (0°C and 10°C). Expression levels increased continuously at 30°C and reached a 2.2-fold maximum compared with the control (20°C) at 12 h (Figure 3), a trend not entirely consistent with the qRT-PCR results.




Figure 2 qRT-PCR analysis of *Hsp845* gene expression in response to temperature stress. Standard error bars are shown: * $P < 0.05$; ** $P < 0.01$.




Figure 3 Western blot analysis of *Hsp845* protein expression in response to temperature stress.

3.3 Expression characteristics of *Hsp845* in response to salinity stress

With regard to salinity stress, *Hsp845* gene expression was

found to be immediately and significantly enhanced by both low salinity (0‰) and high salinity (90‰ and 120‰) during the first 2 h, but was not sensitive to 15‰ salinity (Figure 4). Over the next 10 h, *Hsp845* expression decreased under exposure to salinity levels of 0‰ and 90‰. At a salinity level of 120‰, however, *Hsp845* gene expression reached a maximum 44.8-fold increase compared with the control (45‰ salinity) at 6 h and then exhibited a 9.8-fold decrease over the next 6 h relative to the control. *Hsp845* expression was also enhanced by high salinity (90‰ and 120‰) and reached maximum relative expression increases of 1.9-fold and 1.7-fold after 12 h, respectively. In the absence of salt, *Hsp845* protein expression was significantly inhibited, contrary to the qRT-PCR results. Similarly, *Hsp845* expression was not sensitive to a salinity level of 15‰ (Figure 5).




Figure 4 qRT-PCR analysis of *Hsp845* gene expression in response to salinity stress. Standard error bars are shown: * $P < 0.05$; ** $P < 0.01$.




Figure 5 Western blot analysis of *Hsp845* protein expression in response to salinity stress.

3.4 Expression characteristics of *Hsp845* in response to combined temperature and salinity stress

qRT-PCR analysis revealed that *Hsp845* gene expression was enhanced by heat treatment at 30°C at a salinity of 15‰ or 90‰, but was inhibited at 0°C at those salinity levels. After exposure to 30°C, the transcriptional expression of *Hsp845*

reached a maximum level that was 3.2-fold (under 15‰ salinity) and 4.0-fold (90‰ salinity) higher at 6 h compared with the control (20°C, 45‰ salinity), and then decreased during the next 6 h (Figure 6). The expression characteristics of *Hsp845* in response to combined temperature and salinity stress were similar to those detected by qRT-PCR. High temperature enhanced *Hsp845* protein expression under both low (15‰) and high (90‰) salinity, with maximum increases of 1.7-fold and 1.8-fold respectively observed at 12 h compared with the control (20°C, 45‰ salinity). By contrast, treatment at 0°C with a salinity of 15‰ or 90‰ inhibited *Hsp845* expression (Figure 7).




Figure 6 qRT-PCR analysis of *Hsp845* gene expression in response to combined temperature and salinity stress. Standard error bars are shown: * $P < 0.05$; ** $P < 0.01$.




Figure 7 Western blot analysis of *Hsp845* protein expression in response to combined temperature and salinity stress.

4 Discussion

In this study, we investigated the expression characteristics of the *Hsp845* gene at both transcriptional and translational levels in the Antarctic psychrotrophic bacterium *Psychrobacter* sp. G under various stress treatments using qRT-PCR and western blotting, respectively.

Hsp845, which is affiliated with GrpE, is the co-

chaperone of *Hsp70*^[33]. In this study, high temperature (30°C) significantly enhanced *Hsp845* expression at both transcriptional and translational levels. Previous studies have also indicated that the *Hsp70* chaperone is generally induced in response to an external temperature rise of 5°C–10°C above the optimal growth temperature of an organism^[29] and that GrpE expression is rapidly and transiently induced in response to elevated temperature^[13].

Clark et al.^[30] cloned *Hsp70* family genes *Hsc70*, *Hsp70*, and *Grp78* from Antarctic mollusks *Laternula elliptica* and *Nacella concinna* and studied the expression characteristics of the *Hsp845* gene under high temperatures (15°C and 20°C) by qRT-PCR. When exposed to high temperatures, the expression of *Hsp70* was found to be significantly induced, indicating that this gene plays an important role in the high temperature adaptation of Antarctic mollusks. Park et al.^[31] also studied the expression characteristics of *Hsp70* under high temperature conditions in the Antarctic clam *Laternula elliptica*. They observed that high temperature (10°C) significantly increased *Hsp70* gene expression in gills and digestive gland tissue. The expression of *Hsp70* reached a maximum (4.6-fold greater compared with the control) at 12 h in gill tissue, whereas maximum expression (3.6-fold greater compared with the control) in digestive gland tissue was observed at 24 h. Liu et al.^[22] cloned the 2232-bp full-length sequence of the *Hsp70* gene (*CiHsp70*) of the Antarctic ice alga *Chlamydomonas* sp. ICE-L from a cDNA library. qRT-PCR showed that both cold and heat shock treatments induced *CiHsp70* expression. Under high salinity (62‰) stress, *CiHsp70* expression was induced 3.0-fold compared with the control (31‰). Ultraviolet radiation induced expression of *CiHsp70* gene expression as well. Briefly, *CiHsp70*, as a chaperone, may play an important role in *Chlamydomonas* sp. ICE-L adaption to the polar environment.

Hsp845 protein expression at the translational level lagged behind transcriptional-level expression and reached a maximum at 12 h after exposure to 30°C. The highest relative expression level, 2.2-fold greater than that of the control (20°C), was much lower than that of transcriptional-level expression (5.3-fold greater compared with the control). The same tendency was also observed in the other two stress treatments, indicating that regulation may occur between transcriptional and translational levels.

Freezing and thawing of sea ice during different seasons in Antarctica, the habitat of *Psychrobacter* sp. G, affects the salinity of offshore seawater. Kilstrup et al.^[19] have reported the *Hsp70* chaperone is induced during salinity stress in *Lactococcus lactis*, while Nath et al.^[32] have additionally discovered that hypersalinity creates a more general stress response than hyposaline conditions. The former authors observed that the *Hsp70* chaperone had a transient response after a shift to low or high salinity during adaptation to a particular salinity. This finding may explain why *Hsp845* gene expression was immediately and significantly enhanced at the transcriptional level by both low (0‰) and high (90‰ and 120‰) salinity during the first 2 h of treatment. The effects of

high salinity (120‰) on *Hsp845* gene expression were longer lasting, as the maximum relative expression level was reached at 6 h. In our study, however, we found that the *Hsp845* protein was over-expressed at the transcriptional level after exposure to a salinity of 0‰, but was obviously inhibited at the translational level. The functional contribution of Hsps to salinity stress adaptation is still not clear. One possibility is that Hsps are induced as a consequence of the accumulation of denatured proteins resulting from lowered water activity during plasmolysis of salinity-shocked cells^[33].

Psychrobacter sp. G may simultaneously encounter multiple stress factors. For example, freezing is often accompanied by salinity stress caused by the absence of unfrozen water^[21,34]. We therefore also investigated the combined effect of temperature and salinity on *Hsp845* gene expression. qRT-PCR revealed that *Hsp845* gene expression was enhanced by treatment at 30°C in combination with 15‰ or 90‰ salinity, but was inhibited at 0°C under the same salinity levels. The results of western blotting also uncovered the same trend. Nonetheless, *Hsp845* appeared to be more sensitive to high temperature than to salinity in the combined stress treatment at both transcriptional and translational levels, a observation inconsistent with the results of the single-factor experiment.

In this study, we found that *Hsp845* was induced at both transcriptional and translational levels by high temperature (30°C). *Hsp845* was induced at the transcriptional level by both low (0‰) and high (90‰ and 120‰) salinity, but was only induced at the translational level by high salinity (90‰ and 120‰). In the single-factor experiment, the relative expression of *Hsp845* under salinity stress was far higher than that observed under temperature stress. Under combined stress, however, *Hsp845* was more sensitive to high temperature than to different salinities at both transcriptional and translational levels. The mechanism responsible for this inconsistent pattern needs further study. The maximum relative expression at the transcriptional level was higher than that at the translational level; moreover, the maximum expression of *Hsp845* at the translational level lagged behind that at the transcriptional level. Finally, expression at the transcriptional level in the absence of salt seemed to conflict with results at the translational level; the underlying mechanism also needs to be studied further. Although the mechanism of GrpE proteins contributing to adaption to various stresses in microorganisms living in Antarctic seawater is still far from being understood, we believe that the present work is a good starting point for further investigations.

Acknowledgements This work was supported financially by the National Natural Science Foundation of China (Grant no. 41176174).

References

- Feder M E, Hofmann G E. Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology. *Annu Rev Physiol*, 1999, 61(1): 243–282
- Che S, Song W Z, Lin X Z. Response of heat-shock protein (*HSP*) genes to temperature and salinity stress in the Antarctic psychrotrophic bacterium *Psychrobacter* sp. G. *Curr Microbiol*, 2013, 67(5): 601–608
- Sugimoto S, Saruwatari K, Higashi C, et al. The proper ratio of GrpE to DnaK is important for protein quality control by the DnaK-DnaJ-GrpE chaperone system and for cell division. *Microbiology*, 2008, 154(7): 1876–1885
- McCarty J S, Buchberger A, Reinstein J, et al. The role of ATP in the functional cycle of the DnaK chaperone system. *J Mol Biol*, 1995, 249(1): 126–137
- Russell R, Jordan R, McMacken R. Kinetic characterization of the ATPase cycle of the DnaK molecular chaperone. *Biochemistry*, 1998, 37(2): 596–607
- Szabo A, Langer T, Schröder H, et al. The ATP hydrolysis-dependent reaction cycle of the *Escherichia coli* Hsp70 system DnaK, DnaJ, and GrpE. *Proc Natl Acad Sci U S A*, 1994, 91(22): 10345–10349
- Liberek K, Marszalek J, Ang D, et al. *Escherichia coli* DnaJ and GrpE heat shock proteins jointly stimulate ATPase activity of DnaK. *Proc Natl Acad Sci U S A*, 1991, 88(7): 2874–2878
- Siegenthaler R K, Christen P. The importance of having thermosensor control in the DnaK chaperone system. *J Biol Chem*, 2005, 280(15): 14395–14401
- Brehmer D, Gässler C, Rist W, et al. Influence of GrpE on DnaK-substrate interactions. *J Biol Chem*, 2004, 279(27): 27957–27964
- Zylicz M, Ang D, Georgopoulos C. The grpE protein of *Escherichia coli*. Purification and properties. *J Biol Chem*, 1987, 262(36): 17437–17442
- Yura T, Nagai H, Mori H. Regulation of the heat-shock response in bacteria. *Annu Rev Microbiol*, 1993, 47(1): 321–350
- Harrison C. Grp E, a nucleotide exchange factor for DnaK. *Cell Stress Chaperones*, 2003, 8(3): 218–224
- Lipinska B, King J, Ang D, et al. Sequence analysis and transcriptional regulation of the *Escherichia coli* *grpE* gene, encoding a heat shock protein. *Nucl Acids Res*, 1988, 16(15): 7545–7562
- Wu B, Ang D, Snavely M, et al. Isolation and characterization of point mutations in the *Escherichia coli* *grpE* Heat Shock Gene. *J Bacteriol*, 1994, 176(22): 6965–6973
- Barthel S, Rupperecht E, Schneider D. Thermostability of two cyanobacterial GrpE thermosensors. *Plant Cell Physiol*, 2011, 52(10): 1776–1785
- Bukau B, Walker G C. Cellular defects caused by deletion of the *Escherichia coli* *dnaK* gene indicate roles for heat shock protein in normal metabolism. *J Bacteriol*, 1989, 171(5): 2337–2346
- Hartl F U, Hayer-Hartl M. Molecular chaperones in the cytosol: from nascent chain to folded protein. *Science*, 2002, 295(5561): 1852–1858
- Lindquist S, Craig E A. The heat-shock proteins. *Annu Rev Genet*, 1988, 22(1): 631–677
- Kilstrup M, Jacobsen S, Hammer K, et al. Induction of heat shock proteins DnaK, GroEL, and GroES by salt stress in *Lactococcus lactis*. *Appl Environ Microbiol*, 1997, 63(5): 1826–1837
- Neidhardt F C, Van Bogelen R A. Heat shock response//Neidhardt F C. *Escherichia coli* and *Salmonella typhimurium*: cellular and molecular biology. Washington, DC: American Society for Microbiology, 1987: 1334–1345
- Thomas D N, Dieckmann G S. Antarctic Sea ice—a habitat for extremophiles. *Science*, 2002, 295(5555): 641–644
- Liu S H, Zhang P Y, Cong B L, et al. Molecular cloning and expression analysis of a cytosolic *Hsp70* gene from Antarctic ice algae *Chlamydomonas* sp. ICE-L. *Extremophiles*, 2010, 14(3): 329–337

- 23 Holtmann G, Brigulla M, Steil L, et al. RsbV-independent induction of the SigB-dependent general stress regulon of *Bacillus subtilis* during growth at high temperature. *J Bacteriol*, 2004, 186(18): 6150–6158
- 24 Grimshaw J P A, Jelesarov I, Siegenthaler R K, et al. Thermosensor action of GrpE: the DnaK chaperone system at heat shock temperatures. *J Biol Chem*, 2003, 278(21): 19048–19053
- 25 Lin X Z, Cui S S, Xu G Y, et al. Cloning and heterologous expression of two cold-active lipases from the Antarctic bacterium *Psychrobacter* sp. G. *Polar Res*, 2010, 29(3): 421–429
- 26 Song W Z, Lin X Z, Huang X H. Characterization and expression analysis of three cold shock protein (CSP) genes under different stress conditions in the Antarctic bacterium *Psychrobacter* sp. G. *Polar Biol*, 2012, 35(10): 1515–1524
- 27 Livak K J, Schmittgen T D. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method. *Methods*, 2001, 25(4): 402–408
- 28 Yu S S, Yang H, Chai Y M, et al. Molecular cloning and characterization of a C-type lectin in roughskin sculpin (*Trachidermus fasciatus*). *Fish Shellfish Immunol*, 2013, 34(2): 582–592
- 29 Lindquist S. The heat-shock response. *Annu Rev Biochem*, 1986, 55(1): 1151–1159
- 30 Clark M S, Fraser K P P, Peck L S. Antarctic marine molluscs do have an HSP70 heat shock response. *Cell Stress Chaperones*, 2008, 13(1): 39–49
- 31 Park H, Ahn I Y, Lee H E. Expression of heat shock protein 70 in the thermally stressed antarctic *Clam Laternula elliptica*. *Cell Stress Chaperones*, 2007, 12(3): 275–282
- 32 Nath I V A, Bharathi P A L. Diversity in transcripts and translational pattern of stress proteins in marine extremophiles. *Extremophiles*, 2011, 15(2): 129–153
- 33 Overpeck J, Hughen K, Hardy D, et al. Arctic environmental change of the last four centuries. *Science*, 1997, 278(5341): 1251–1256
- 34 Pearce D A. Climate change and the microbiology of the Antarctic Peninsula region. *Sci Prog*, 2008, 91(2): 203–217